

# Jurnal 1 Polyglactin 910 (Vicryl Rapide)

*by Admin Perpustakaan Stikesyo*

---

**Submission date:** 30-Mar-2023 03:28AM (UTC-0400)

**Submission ID:** 2050787795

**File name:** jurnal\_amilya.pdf (49.89K)

**Word count:** 2239

**Character count:** 11455

Research Article

9  
**Polyglactin 910 (Vicryl Rapide) for Reducing Pain and Wounds Healing in Postpartum Perineum**

***Polyglactin 910 (Vicryl Rapide) untuk Mengurangi Nyeri dan Mempercepat Penyembuhan Luka pada Perineum Postpartum***

**Amilya Pradita, Abkar Raden, Farida Kartini**

*Faculty of Health Science Universitas Aisyiyah  
Yogyakarta*

**Abstract**

**Objective:** The purpose of this study was to compare the use of polyglactin 910 (vicrylrapide) and chromic catgut to perineal pain and healing wounds on postpartum.

**Methods:** This research is true experiment with a randomized controlled trial (RCT) and single-blind. The number of respondents pad this study of 40 respondents were divided into 20 respondents to the intervention group (vicrylrapide) and 20 respondents for the control group (chromic catgut). Data analysis using independent t-test.

**Result:** Statistical test results using independent t-test for pain showed  $p = 0.035 < 0.05$  and for wound healing showed  $p = 0.000 < 0.05$  of a second can mean the results are there differences in the use of threads polyglactin 910 (vicrylrapide) and chromic catgut thread to perineal pain and healing wounds in the mother postpartum perineum.

**Conclusions:** The use rapidevicryl thread to repair perineal perineum or stitches can reduce perineal pain and healing wounds better than the use of the chromic catgut thread.

**Keywords:** chromic catgut, pain, polyglactin 910, rapide vicryl, wound healing of the perineum.

**Abstrak**

**Tujuan:** Untuk mengetahui perbandingan penggunaan polyglactin 910 (vicryl rapide) dan chromic catgut terhadap nyeri dan penyembuhan luka perineum pada postpartum.

**Metode:** Penelitian ini merupakan penelitian true experiment dengan randomized controlled trial (RCT) dansingle blind. Jumlah responden pada penelitian ini sebanyak 40 responden yang di bagi menjadi 20 responden untuk kelompok intervensi (vicryl rapide) dan 20 responden untuk kelompok kontrol (chromic catgut ). Analisis data menggunakan independent t-test.

**Hasil:** Uji statistik menggunakan t-test independent untuk nyeri didapatkan hasil  $p=0,035<0,05$  dan untuk penyembuhan luka didapatkan hasil  $p=0,000<0,05$  dari kedua hasil tersebut dapat diartikan ada perbedaan penggunaan benang polyglactin 910 (vicryl rapide) dan benang chromic catgut terhadap nyeri perineum dan penyembuhan luka perineum pada ibu postpartum.

**Kesimpulan:** TPenggunaan benang vicryl rapide untuk perbaikan perineum atau penjahitan luka perineum dapat mengurangi nyeri perineum dan penyembuhan luka yang lebih baik dibandingkan dengan penggunaan pada benang chromik catgut.

**Kata kunci:** chromik catgut, nyeri, penyembuhan luka perineum, polyglactin 910, vicryl rapide.

**Correspondence author.** Amilya Pradita. Faculty of Health Science Universitas Aisyiyah Yogyakarta. amilyapradita24@gmail.com

Received: March, 2019 Accepted: March, 2021 Published: July, 2021

## INTRODUCTION

Women who give birth vaginally will undergo perineal laceration / trauma well as spontaneous tearing during childbirth physiological or surgical incision (episiotomy). Spontaneous rupture perineum perineal injuries that occur due to certain reasons without tearing or deliberate action. These injuries occur during delivery and usually irregular<sup>1</sup>. Perineum rupture is one of the trauma that often affects women during childbirth. Perineal trauma and perineal repair is associated with puerperal resulting in morbidity such as bleeding, infection, pain, urinary and faecal incontinence and sexual dysfunction<sup>2</sup>. Other factors that can increase morbidity Other types of materials or stitches are used during repair of perineal<sup>3</sup>.

Suture material for perineal repair used can increase perineal pain in the short-term post-childbirth, thus required the use of suturing best to reduce perineal pain and repair of perineal trauma, so that will reduce the impact of short-term<sup>4</sup>.

Suture material for perineal repair is often used is chromic catgut. Chromic catgut is yarn types absorption (absorption), have tensile strength 17-21 days, 90-110 days old absorption<sup>5</sup>. Other thread types that are rarely used is vicryl rapide is a type of yarn absorption (absorption), has a tensile strength of 10-14 days, 42 days longer absorption<sup>6</sup>.

The use of suture materials are superior in addition to reducing perineal pain also can accelerate wound healing, such as research conducted by explains that the use of polyglactin 910 suture material (vicryl rapide) is an ideal suture material for perineal repair perineal pain that produces fewer and better wound healing compared to the use of chromic catgut suture material<sup>3</sup>.

Based on the above researchers are searching for related to "Comparative Use of polyglactin 910 (Vicryl Rapide) and Chromic catgut Against Pain and Wound Healing perineum on Postpartum"

## METHODS

The research is true experiment (randomized controlled trial) with a single blinding technique.

The population in this study were all postpartum mothers who birth at health centers and health centers Godean I Mlati II in Sleman. The sample in this study is a normal birth mothers who suffered wounds perineum and perineal suturing was done using 910 threads polyglactin (vicryl rapide) or chromic catgut. Sampling was done by consecutive sampling. The sample size in this study were 40 respondents (20 respondents intervention group, the control group of 20 respondents) with the inclusion criteria primigravida pregnant women who spontaneously vaginal delivery without complications or complications, Maternal with spontaneous rupture of 2nd degree perineal and given lidocaine when suturing and mothers who do not smoke. This study were followed up for 15 days to see perineal wound pain and wound healing of the perineum. Measuring tool to see perineal pain using the NRS and wound healing using REEDA.

## RESULTS

This research was conducted in two health centers in Sleman namely Puskesmas Puskesmas Mlati Godean I and II. The number of samples in this study were 40 respondents (20 respondents 20 respondents intervention group and the control group).

**Table 1.** Characteristics of Research Subjects

Characteristics of Respondents	Intervention group (N = 20), %	Control group (N = 20) %
<b>Age (y.o)</b>		
Not at risk		
20-35	19 (95.0)	19 (95.0)
<b>Risk</b>		
(<20 and> 35 )	1 (5.0)	1 (5.0)
Total	20 (100)	20 (100)
<b>Last hemoglobin</b>		
Normal	18 (90.0)	16 (80.0)
mild anemia	2 (10.0)	4 (20.0)
Total	20 (100)	20 (100)
<b>Techniques suturing</b>		
simple interrupted	5 (25.0)	5 (25.0)
simple interrupted and		
simple continuous	15 (75.0)	15 (75.0)
Total	20 (100)	20 (100)
<b>Personal hygiene</b>		
SOP Godean I		
Community Health Center	10 (50.0)	10 (50.0)
SOP PHC Mlati II	10 (50.0)	10 (50.0)
Total	20 (100)	20 (100)

**Table 2.** The Mean Perineal Pain and Healing Wounds

Variables	Days to-	Group	
		Intervention Average	Control Average
Wound pain perineum	1	4.15	5.30
	2	4.00	4.75
	3	3.55	4.05
	4	2.90	3.65
	5	2.55	3.10
	6	2.00	2.60
	7	1.65	2.40
	8	1.40	2.05
	9	1.15	1.50
	10	0.70	1.35
	11	0.50	1.15
	12	0.30	0.70
	13	0.15	0.35
	14	0.05	0.15
	15	0.05	0.15
Perineum Wound Healing	1	4.75	5.65
	3	1.65	2.30
	5	0.35	1.90
	7	0,00	1.75
	9	0,00	0.20
	11	0,00	0.00
13	0,00	0.00	
15	0,00	0.00	

**Table 3.** Comparison of Pain and Wound Healing Perineum Perineum

	Group		Mean Difference	CI 95%	P-value
	Intervention mean $\pm$ SD	Control mean $\pm$ SD			
The perineum pain	25.10 $\pm$ 11.253	33.25 12,311	-8.150	-15.700 – 0.600	0.035
Perineum Wound Healing	6.75 $\pm$ 1.333	11.30 $\pm$ 1.455	-4.550	-5.443 - 3.657	0.000

Results of research on Comparative Use of polyglactin 910 (Vicryl rapide) and Chromic catgut against Pain and Wound Healing perineum in Puskesmas Regency Sleman, it can be concluded use of polyglactin 910 (vicryl rapide) has a better effect on pain and wound healing of the perineum compared with the use of chromic catgut.

### DISCUSSION

Based on the above table 1 showed that the characteristics of the respondents in the intervention group and controls by age in mind that almost all respondents are in the productive age and only 2 respondents who have a risky age. Respondents who experienced mild anemia in the intervention group as much as 2 respondents and in the control group as much

as 4 respondents. Sewing techniques used for engineering disconnected in the intervention group and by 5 respondents in the control group by 5 respondents in the two health centers had more frequent use sewing techniques baste and disconnected. Comparison of the appropriate personal hygiene at the health center respondents SOP intervention group and the control group is the same, each of the 20 respondents according to SOP I godean health centers and 20 health centers of respondents according to SOP Mlati II.

Based on Table 2 above showed a mean value of the highest perineal pain scores in the intervention group, namely on day 1 with a mean value of 4.15 and the lowest on day 14 with a mean value of 0.05, while the highest in the control group to-1 with a mean value of 5.30 and the lowest on day -14 with a mean value of 0.15.

The results mean value for perineal wound healing in the intervention group highest on day 1 with a mean value of 4.75 and the lowest at day-7 is 0.00. Whereas in the control group the highest average value on day 1 with a mean value of 5.65 and the lowest on day 11 with a mean value of 0.00.

Based on Table 3. The test results independent t-test can be concluded that perineal pain and healing wounds in the intervention group mean values of 25.10 and 6.75, whereas in the control group mean values of 33.25 and 11.30. The test results using statistical tests obtained independent t-test p-value <0.05, which means that there are differences in the use of threads polyglactin 910 (vicryl rapide) and chromic catgut thread to perineal pain and healing wounds in the mother postpartum perineum.

This is according to research conducted by explains that the use of suture material with polyglactin 910 (Vicryl rapide) superior than chromic catgut. polyglactin 910 (Vicryl rapide) can be absorbed naturally and may reduce pain compared with the use of chromic catgut sutures<sup>7</sup>. Also in line with research explained that the use of polyglactin 910 suture material (vicryl rapide) is an ideal suture material for perineal repair perineal pain that produces fewer and better wound healing compared to the use of chromic catgut suture material<sup>8</sup>.

Explains that the use of thread vicryl rapide produces less pain and better wound healing (p <0.05 significant)<sup>9</sup>. In concluded that vicryl rapide better at reducing pain and wound healing in wound repair perineum.

Explained that the use of vicryl suture materials rapide there is a significant reduction in short-term pain<sup>10</sup>. The pain began to improve from the third day onwards, no experience moderate pain on day 7 and on day 15, no woman who complained of pain in the group vicryl rapide compared with chromic catgut were still experiencing mild pain on day 15. From day 7 onwards perception of pain was lower in the group vicryl rapide compared with chromic catgut

## CONCLUSION

All in all, vicryl rapide thread to repair perineum or perineal wound suture is more effective to reduce perineal pain and has better perineal wound healing than chromic catgut thread. Health workers are suggested to choose the right material of thread in perineal repair in order to reduce pain and speed up the process of perineal wound healing in postpartum mothers with perineal rupture

## REFERENCES

1. Abu Seada, M. R., Borg, T. F., & Samy, M. M. Continuous versus Interrupted Suturing in Repair of Lateral and Mediolateral Episiotomy: A Randomized Controlled Trial. *Egy J Hos Med*. 2018; 71: 2667–80.
2. Gray, E. Suturing versus non-suturing of spontaneous perineal tears following normal birth: Factors that influence the practice of New Zealand / Aotearoa midwives. 2010:10.
3. Bharathi, A., Dharma Reddy, DB, & Sharath Kote, GS. A prospective randomized comparative study of vicryl rapide versus chromic catgut for episiotomy repair. *J Clin Diagnos Research*. 2013; 7: 326-30.
4. Kettle C, D. T. Continuous and interrupted suturing techniques for repair of episiotomy or second-degree tears. *The Cochrane Database of Systematic Reviews*. 2012; 11: CD000947.
5. Dudley, L., Kettle, C., & Ismail, K. Prevalence, pathophysiology and current management of dehiscent perineal wounds following childbirth. *Bri J Midwifery*. 2013; 21: 160–71.
6. Abdullah, M., Noreen, A., Iqbal, M., & Sohail, R. Comparison Between Chromic Catgut and Vicryl Rapide for Analgesia Requirement in episiotomy Repair in Primigravidas. 2015; 21: 193-6.
7. Samant, M., Bose, E., Ghosh, A. Lal, P., & Mishra, S. Comparison of the impact of polyglactin 910 (Vicryl rapide) and chromic catgut sutures on perineal pain following episiotomy wound repair in the eastern Indian Patients. *J Scien Soc*. 2013; 40: 95.
8. Perumal, D., & Selvaraju, D. Comparative study of episiotomy repair: versus synthetic absorbable chromic catgut suture material. *Int J Reprod Contracept Obstet Gynecol*. 2017; 6: 2186.
9. Esa B., M. Samant, P. Lal, S. Mishra, Ghost A. Comparison of the impact of polyglactin 910 (Vicrylrapide) and chromic catgut sutures on perineal pain following episiotomy wound repair in the eastern Indian Patients. *J Scien Soc*. 2018; 40.
10. Odijk, R., Hennipman, B., Rousian, M., Madani, K., Dijksterhuis, M., de Leeuw, J. W., & van Hof, A. The MOVE-trial: Monocryl® vs. Vicryl rapide™ for skin repair in mediolateral episiotomies: A randomized controlled trial. *BMC Pregnancy and Childbirth*. 2017; 17: 1–7.

# Jurnal 1 Polyglactin 910 (Vicryl Rapide)

---

## ORIGINALITY REPORT

---

12%

SIMILARITY INDEX

7%

INTERNET SOURCES

9%

PUBLICATIONS

1%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- 1** Chandrakant S Madkar, Hemant Deshpande, Apurva Singh, Akanksha Allahbadia. "Prospective randomized control trial comparing cromic catgut 1-0 versus fast absorbing polyglactin 910 2-0 for episiotomy repair in a semi-urban Indian population group", Indian Journal of Obstetrics and Gynecology Research, 2019  
Publication 2%

---
- 2** Ardiana Batubara, Fitriyani Pulungan. "The Effect of Pineapple Juice (Ananas comosus) on the Uterine Involution of Postpartum Mothers at Pratama Jannah and Dermawati Maternity Clinics of Deli Serdang Regency in 2019", Global Journal of Health Science, 2022  
Publication 1%

---
- 3** [www.scirp.org](http://www.scirp.org)  
Internet Source 1%

---
- 4** Firdaus Ujang, Rosnah Sutan. "The Effects of Short Health Messages Intervention in Improving Knowledge and Attitude on Sexual 1%

and Reproductive Health among Late Adolescents at Colleges in Malaysia", Open Journal of Preventive Medicine, 2018

Publication

5

C Kettle. "Continuous versus interrupted sutures for perineal repair", Cochrane Database of Systematic Reviews Reviews, 01/26/1998

Publication

1 %

6

[www.ijrcog.org](http://www.ijrcog.org)

Internet Source

1 %

7

Submitted to Higher Education Commission Pakistan

Student Paper

1 %

8

Carolina López-Lapeyrere, Montserrat Solís-Muñoz, Ana Belén Hernández-López, Ricardo Rodríguez-Barrientos et al. "Perineal repair of media-lateral episiotomies and 2nd degree tears by midwives: A randomised controlled trial comparing three suture techniques", International Journal of Nursing Studies, 2020

Publication

1 %

9

[discovery.researcher.life](http://discovery.researcher.life)

Internet Source

1 %

10

Ringdal, G.I.. "Family Satisfaction with End-of-Life Care for Cancer Patients in a Cluster

1 %

Randomized Trial", Journal of Pain and Symptom Management, 200207

Publication

11

hub.op.ac.nz

Internet Source

1 %

12

www.jscisociety.com

Internet Source

1 %

13

Arianti Arianti, Nadila Putri Mayna, Yuda Hidayat. "MOBILISASI DINI TERHADAP PEMULIHAN PERISTALTIK USUS DAN SKALA NYERI PASIEN POST PEMBEDAHAN", Journal of Holistic Nursing Science, 2020

Publication

1 %

14

www.tandfonline.com

Internet Source

1 %

15

Ni Wayan Rahayu Ningtyas, RR Sri Endang Pujiastuti, Nina Indriyawati. "EFFECTIVENESS OF PROGRESSIVE MOBILIZATION LEVEL I AND II ON HEMODYNAMIC STATUS AND DECUBITUS ULCER RISK IN CRITICALLY ILL PATIENTS", Belitung Nursing Journal, 2017

Publication

1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On



# Jurnal 1 Polyglactin 910 (Vicryl Rapide)

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---